MA 4207 - Mathematical Logic
Course-Webpage `http://www.comp.nus.edu.sg/~fstephan/mathlogicug.html`
Homework

Frank Stephan. Departments of Mathematics and Computer Science,
10 Lower Kent Ridge Road, S17#07-04 and 13 Computing Drive, COM2#03-11,
National University of Singapore, Singapore 119076.
Email fstephan@comp.nus.edu.sg
Telephone office 65162759 and 65164246
Office hours Thursday 14.00-15.00h at Mathematics S17#07-04

Homework 11.1
Assume that α, β, γ are well-formed formulas. Give a formal proof of the statement

$$\{\beta, \gamma\} \models \alpha \to \beta$$

which only uses the formulas from Λ and the Modus Ponens.

Homework 11.2
Assume that $\{\alpha, \beta\}$ tautologically implies γ. The below derivation is incorrect. Say what the fault is and replace it by a corrected one:

1. $\{\alpha, \beta\} \vdash \alpha \to \beta \to \gamma$ (Axiom Group 1)
2. $\{\alpha, \beta\} \vdash \beta$ (Copy)
3. $\{\alpha, \beta\} \vdash \beta \to \alpha \to \beta$ (Axiom Group 1)
4. $\{\alpha, \beta\} \vdash \alpha \to \beta$ (Modus Ponens)
5. $\{\alpha, \beta\} \vdash \gamma$ (Modus Ponens)

For the following exercises, P, Q are predicates and a, b, c are constants.

Homework 11.3
Make a formal proof for

$$\{\forall x\, [P(x) \to Q(c)], \forall x\, [\neg P(x) \to Q(c)]\} \vdash Q(c)$$

Homework 11.4
Make a formal proof for $\{\forall x\, [P(x)], \exists y\, [\neg P(y)]\} \vdash Q(z)$.

Homework 11.5
Make a formal proof for $\emptyset \vdash \forall x\, \forall y\, [P(x) \to Q(y)] \to P(a) \to Q(b)$.

Homework 11.6
Is the statement $\emptyset \vdash P(x) \to \forall y\, [P(y)]$ correct? Explain your answer.

Homework 11.7
Is the statement $\emptyset \vdash P(x) \to \forall y\, [P(x)]$ correct? Explain your answer.

Homework 11.8
Is the statement $\emptyset \vdash P(x) \to \exists y\,[P(y)]$ correct? Explain your answer.

Homework 11.9
Let $(G, \circ, f, e)$ be a structure and Γ contain the following axioms:

- $\forall x, y, z\,[(x \circ y) \circ z = x \circ (y \circ z)]$;
- $\forall x, y\,[x \circ y = y \circ x]$;
- $\forall x\,[x \circ e = x]$;
- $\forall x\,[x \circ f(x) = e]$;
- $\forall x, y, z\,[x \circ y = x \circ z \to y = z]$;

So $(G, \circ)$ is an Abelian group with neutral element e and inversion f. Prove informally the following results:

- $\forall v, w\,[f(v) = f(w) \to v = w]$;
- $\forall v, w\,[v \circ w = e \to f(v) = w]$;
- $\forall v, w\,[f(v \circ w) = f(w) \circ f(v)]$.

Homework 11.10
Consider all structures $(A, \circ)$ where A has two elements and satisfies the axioms

$$\forall x\,[x \circ x = x] \text{ and } \forall x\,\forall y\,[x \circ y = y \circ x].$$

Show that all these structures are isomorphic.

Homework 11.11
Assume that $(\mathbb{N}, +, <, 0, 1, P)$ is a structure where $\mathbb{N}$ is the set of natural numbers and $+, <, 0, 1$ have the usual meaning on $\mathbb{N}$. Let the powers of 2 be the set $\{1, 2, 4, 8, 16, \ldots\}$ and make a formula α such that $(\mathbb{N}, +, <, 0, 1, P) \models \alpha$ iff $\forall x\,[Px \leftrightarrow \text{x is a power of 2}]$.

Note that such a formula only implicitly defines the powers of 2 and not explicitly; therefore this formula α does *not say* that the powers are definable from addition and order in $\mathbb{N}$.

Homework 11.12
Make a formula α which says that $f : A \to A$ is a one-to-one function but not an onto function. Provide a model $(A, f, =)$ which satisfies α. Can A be finite?